



Wind field estimation via C^1 Bézier smoothing on manifolds

Pierre-Yves Gousenbourger, Estelle Massart
`pierre-yves.gousenbourger@uclouvain.be`

Université catholique de Louvain, ICTEAM


13 décembre 2018









Poké-Collab: Kanto / 151 Pokemon by 151 Artists
 July 22 - August 10 2013



The wind field estimation






How to fit a curve to points on \mathcal{M} ?

1D : Interpolative Bézier curves


Each segment between two consecutive points is a **Bézier curve** of degree K .




$$\beta_K(t, \mathbf{b}) = \sum_{i=0}^K b_i B_{iK}(t)$$

[G. et al. 2014, Arnould et al. 2015]

Reconstruction : the De Casteljau algorithm




The straight line is a geodesic



(


The exponential map to construct the geodesic

$$\gamma(t) = \text{Exp}_x(t\xi_x)$$




The logarithmic map to determine the starting velocity

$$\text{Log}_x(y) = \xi_x$$






Example on the sphere



It's ugly. Make it **smooth**!

Well?! What about fitting, now?


Smooth fitting with Bézier (in \mathbb{R}^n)



Each segment is a Bézier curve smoothly connected!

Unknowns : b_i^- , b_i^+ , p_i .

Differentiability



$$p_i = \frac{b_i^- + b_i^+}{2}$$

Optimal \mathcal{C}^1 -piecewise Bézier fitting (in \mathbb{R}^n)

Minimization of the mean squared acceleration of the path

$$\min_{p_0, b_i^-, b_i^+, p_n} \int_0^1 \|\ddot{\beta}_2^0\|^2 dt + \sum_{i=1}^{n-1} \int_0^1 \|\ddot{\beta}_3^i\|^2 dt + \int_0^1 \|\ddot{\beta}_2^n\|^2 dt + \lambda \sum_{i=0}^n \|d_i - p_i\|_2^2$$

$$\min_{p_0, b_i^-, b_i^+, p_n} \int_0^1 \|\ddot{\beta}_2^0\|^2 dt + \sum_{i=1}^{n-1} \int_0^1 \|\ddot{\beta}_3^i\|^2 dt + \int_0^1 \|\ddot{\beta}_2^n\|^2 dt + \lambda \sum_{i=0}^n \|d_i - p_i\|_2^2$$

Second order polynomial $P(p_0, b_i^-, b_i^+, p_n, \lambda)$

$$\min_{p_0, b_i^-, b_i^+, p_n} \int_0^1 \|\ddot{\beta}_2^0\|^2 dt + \sum_{i=1}^{n-1} \int_0^1 \|\ddot{\beta}_3^i\|^2 dt + \int_0^1 \|\ddot{\beta}_2^n\|^2 dt + \lambda \sum_{i=0}^n \|d_i - p_i\|_2^2$$

Second order polynomial $P(p_0, b_i^-, b_i^+, p_n, \lambda)$

$$\nabla P(p_0, b_i^-, b_i^+, p_n, \lambda)$$

Optimal \mathcal{C}^1 -piecewise Bézier fitting (on \mathcal{M})

- The control points are given by :

$$x_i = \sum_{j=0}^n q_{i,j}(\lambda) d_j$$

- These points are invariant under translation, *i.e.* :


$$x_i - d^{ref} = \sum_{j=0}^n q_{i,j}(\lambda) (d_j - d^{ref})$$

- On manifolds : projection to the **tangent space** of d^{ref} with the **Log**, as $a - b \Leftrightarrow \text{Log}_b(a)$

$$v_i = \text{Log}_{d^{ref}}(x_i) = \sum_{j=0}^n q_{i,j}(\lambda) \text{Log}_{d^{ref}}(d_j)$$

- Back to the manifold with the **Exp** : $x_i = \text{Exp}_{d^{ref}}(v_i)$, where $d^{ref} = d_i$ if x_i is b_i^- , p_i , b_i^+ .

Application : Wind field estimation on $S_+(r, p)$.



Application : Wind field estimation on $S_+(r, p)$.

How to estimate the error ?

- Test set : $\{C(\theta_i)\}_{i \in I_T}$ $I_T = \{1, 3, \dots, 33\}$;
Validation set : $\{C(\theta_i)\}_{i \in I_V}$ $I_V = \{2, 4, \dots, 32\}$;


- Bézier spline $\mathbf{B}(\theta)$ with input data points from I_T

- Mean Squared Error :

$$\text{MSE}(\mathbf{B}(\theta)) = 10 \log \left(\frac{\sum_{i \in I_\Omega} \|C(\theta_i) - \mathbf{B}(\theta_i)\|_F^2}{\sum_{i \in I_\Omega} \|C(\theta_i)\|_F^2} \right), \quad \Omega = \{I, V\}.$$


Application : Wind field estimation on $S_+(r, p)$.

No noise on data



Application : Wind field estimation on $S_+(r, p)$.

With artificial noise (8dB) on data



Fitting with Bézier : pros and cons

✓ Optimality conditions are a closed form linear system.

✓ Method only needs exp and log maps.

✓ The curve is \mathcal{C}^1 .

✗ No guarantee on the optimality when \mathcal{M} is not flat.

✓ We can do denoising.

Paper submitted at the ESANN conference, 2017. Joint work with MIT.

Conclusions

General C^1 -**interpolative/fitting** methods on **manifolds**...
with applications in medical imaging, wind estimation, model
reduction,...


light • closed form • uses few elements in \mathcal{M}

Summary on interpolation :

“Differentiable Piecewise-Bézier Surfaces on Riemannian Manifolds”

[Absil, Gousenbourger, Striewski, Wirth, *SIAM Journal on Imaging Sciences*,
to appear].

Any questions ?



Wind field estimation via C^1 Bézier smoothing on manifolds

Pierre-Yves Gousenbourger, Estelle Massart
pierre-yves.gousenbourger@uclouvain.be

Université catholique de Louvain, ICTEAM

13 décembre 2018